# The Canadian Entomologist

**Index for Volume 137 (2005) / Index pour le volume 137 (2005)**

**Entomological Society of Canada /**
**Société d'entomologie du Canada**

© 2005 Entomological Society of Canada

Athanassiou CG. See Kavallieratos *et al.*, 516
Barber KN. See Pearce *et al.*, 233
Barclay HJ, Li C, Benson L, Taylor S, Shore T. Effects of fire return rates on traversability of lodgepole pine forests for mountain pine beetle (Coleoptera: Scolytidae) and the use of patch metrics to estimate traversability, 566
Bates SL. Damage to common plumbing materials caused by overwintering *Leptoglossus occidentalis* (Hemiptera: Coreidae), 492
Battisti A. See Rouault *et al.*, 450
Baur H. See Gibson *et al.*, 381
Beguin J. See Buddle *et al.*, 120
Behan-Pelletier VM, Eamer B. *Zachvatkinibates* (Acari: Oribatida: Mycobatidae) of North America, with descriptions of sexually dimorphic species, 631
Benson L. See Barclay *et al.*, 566
Bergh JC. See Short and Bergh, 67
Biron DG, Coderre D, Fournet S, Nénon JP, Le Lannic J, Boivin G. Larval respiratory systems of two anthomyiid flies, *Delia radicum* and *Delia antiqua* (Diptera: Anthomyiidae), 163
Boiteau G. Role of vision throughout the flight initiation sequence of the Colorado potato beetle, 420
Boivin G. See Biron *et al.*, 163
See Cournoyer and Boivin, 230
Bolduc E. See Buddle *et al.*, 120
Bologna MA. See Di Giulio *et al.*, 657
Bondrup-Nielsen S. See Teichert and Bondrup-Nielsen, 192
Borden JH. See Broberg *et al.*, 716
See Campbell and Borden, 719
Borys RR. See Fettig *et al.*, 748
Brandl D. See Kuenen *et al.*, 117
Broberg CL, Borden JH, Gries R. Antennae of *Cryptorhynchus lapathi* (Coleoptera: Curculionidae) detect two pheromone components of coniferophagous bark beetles in the stems of *Salix sitchensis* and *Salix scouleriana* (Salicaceae), 716
Brodeur J. See Meloche *et al.*, 226
Buck M. Two introduced spider wasps (Hymenoptera: Pompilidae) new to Canada, with notes on nesting habits and the incidence of introductions, 278
See Marshall *et al.*, 416
Buddle CM. See Lessard and Buddle, 215
Buddle CM, Beguin J, Bolduc E, Mercado A, Sackett TE, Selby RD, Varady-Szabo H, Zeran RM. The importance and use of taxon sampling curves for comparative biodiversity research with forest arthropod assemblages, 120
Burgess LW. See Conway *et al.*, 622
Burton KB. See Conway *et al.*, 622
Butrón A, Ordás B, Revilla P, Sandoya G, Ordás A, Malvar RA. Is leaf or sheath antibiosis involved in the resistance of maize composite EPS12 to *Sesamia nonagrioides*?, 350
Campbell SA, Borden JH. Bark reflectance spectra of conifers and angiosperms: implications for host discrimination by coniferophagous bark and timber beetles, 719
Cantini R. See Rouault *et al.*, 450
Cárcamo HA, Dosdall LM, Johnson D, Olfert O. Evaluation of foliar and seed treatments for control of the cabbage seedpod weevil (Coleoptera: Curculionidae) in canola, 476
Chaoliang L. See Xinhua *et al.*, 83
Cluck DR. See Fettig *et al.*, 748
Coderre D. See Biron *et al.*, 163
Conway HE, Burton KB, Hendrix CA, Burgess LW, Culin JD. Comparison of two different box styles for mass rearing of *Sasajiscymnus tsugae* (Coleoptera: Coccinellidae), a biological control agent of hemlock woolly adelgid (Hemiptera: Adelgidae), 622
Cook J. See Peck and Cook, 283
Cournoyer M, Boivin G. Evidence for kairomones used by the egg parasitoid *Anaphes victus* (Hymenoptera: Mymaridae) when searching for its host, 230
Culin JD. See Conway *et al.*, 622
Dabney CP. See Fettig *et al.*, 748
Dai W, Zhang Y, Hu J. Review of the Chinese leafhopper genus *Paralaevicephalus* (Hemiptera: Cicadellidae: Deltocephalinae) with descriptions of two new species from China, 404
Danyk T, Mackauer M, Johnson DL. Reduced food consumption in the grasshopper *Melanoplus sanguinipes* (Orthoptera: Acrididae) parasitized by *Blaesoxipha atlanis* (Diptera: Sarcophagidae), 356

© 2005 Entomological Society of Canada

De Marmels J, Garrison RW. Review of the genus *Leptagrion* in Venezuela with new synonymies and descriptions of a new genus, *Bromeliagrion*, and a new species, *B. rehni* (Zygoptera: Coenagrionidae), 257
Di Giulio A, Pinto JD, Bologna MA. Larval morphology of the Neotropical genus *Epispasta* (Coleoptera, Meloidae), 657
Dosdall LM. See Gibson *et al.*, 381
See Cárcamo *et al.*, 476
See Laffin *et al.*, 672
Du Y, Roe AD, Sperling FAH. Phylogenetic framework for *Dioryctria* (Lepidoptera: Pyralidae: Phycitinae) based on combined analysis of mitochondrial DNA and morphology, 685
Eamer B. See Behan-Pelletier and Eamer, 631
Eberhard WG. Threading a needle with reinforced thread: intromission in *Ceratitis capitata* (Diptera, Tephritidae), 174
Sexual morphology of male *Sepsis cynipsea* (Diptera: Sepsidae): lack of support for lock-and-key and sexually antagonistic morphological coevolution hypotheses, 551
Elliot BG. See Wise *et al.*, 598
El-Sayed AM. See Trimble and El-Sayed, 110
Erbilgin N, Storer AJ, Wood DL, Gordon TR. Colonization of cut branches of five coniferous hosts of the pitch canker fungus by *Pityophthorus* spp. (Coleoptera: Scolytidae) in central, coastal California, 337
Evenden ML. Potential for combining sex pheromones for the forest tent caterpillar (Lepidoptera: Lasiocampidae) and the large aspen tortrix (Lepidoptera: Tortricidae) within monitoring traps targeting both species, 615
Fasseas C. See Kavallieratos *et al.*, 516
Febria CM, Magnusson AK, Williams DD. Seasonal abundance and prey selection of the nymphs of three sympatric species of *Sympetrum* (Odonata: Libellulidae) in an intermittent pond, 723
Fettig CJ, Borys RR, Dabney CP, McKelvey SR, Cluck DR, Smith SL. Disruption of red turpentine beetle attraction to baited traps by the addition of California fivespined ips pheromone components, 748
Fournet S. See Biron *et al.*, 163
García Prieto F, Nieto Nafría JM. Trois nouvelles espèces du genre *Aphis* de l'Espagne, 49
Garrison RW. See De Marmels and Garrison, 257
Garrison RW, von Ellenrieder N. *Neuragrion mysticum* (Odonata: Megapodagrionidae) demystified, 169
Gibson GAP, Baur H, Ulmer B, Dosdall L, Muller F. On the misidentification of chalcid (Hymenoptera: Chalcidoidea) parasitoids of the cabbage seedpod weevil (Coleoptera: Curculionidae) in North America, 381
Gómez-Anaya JA. See Novelo-Gutiérrez and Gómez-Anaya, 61
Gordon TR. See Erbilgin *et al.*, 337
Greaves G. See Gries *et al.*, 471
Greaves M. See Gries *et al.*, 471
Gries R. See Broberg *et al.*, 716
Gries R, Schaefer PW, Yoo HJS, Greaves M, Gries G. (*Z,E*-6,8-heneicosadien-11-one: major sex-pheromone component of *Orgyia vetusta* (Lepidoptera: Lymantriidae), 471
Grieshop MJ. See Nansen *et al.*, 202
Gut LJ. See Stelinski *et al.*, 98
Hanula JL. See Ulyshen *et al.*, 251
Hemachandra KS, Holliday NJ, Klimaszewski J, Mason PG, Kuhlmann U. Erroneous records of *Aleochara bipustulata* from North America: an assessment of the evidence, 182
Hendrix CA. See Conway *et al.*, 622
Holliday NJ. See Wylie *et al.*, 91
See Hemachandra *et al.*, 182
Hoover K. See Morewood *et al.*, 376
Horn S. See Ulyshen *et al.*, 251
Horton DR, Lewis TM, Neven LG. Ovarian development and lipid reserves are affected by mating delays in three species of *Anthocoris* (Hemiptera: Anthocoridae), 328
Howden HF, Malý V. *Cryptoryctes tricornutus* sp. nov. from the Northern Territory, Australia (Scarabaeidae, Dynastinae, Oryctini), 274
Hu J. See Dai *et al.*, 404
Huang M. See Zhang and Huang, 410
Jayas DS. See Jian *et al.*, 71

© 2005 Entomological Society of Canada

Jian F, Jayas DS, White NDG. Effects of temperature acclimation and age on movement of *Cryptolestes ferrugineus* (Coleoptera: Laemophloeidae) adults in response to temperature gradients, 71
Johnson DL. See Danyk *et al.*, 356
See Cárcamo *et al.*, 476
Johnson ML. See Nansen *et al.*, 202
Kavallieratos NG, Tomanović Ž, Athanassiou CG, Starý P, Žikić V, Sarlis GP, Fasseas C. Aphid parasitoids infesting cotton, citrus, tobacco, and cereal crops in southeastern Europe: aphid–plant associations and keys, 516
Klimaszewski J. See Hemachandra *et al.*, 182
Klimaszewski J, Sweeney J, Price J, Pelletier G. Rove beetles (Coleoptera: Staphylinidae) in red spruce stands, eastern Canada: diversity, abundance, and descriptions of new species, 1
Knight AL, Light DM. Timing of egg hatch by early-season codling moth (Lepidoptera: Tortricidae) predicted by moth catch in pear ester- and codlemone-baited traps, 728
Developing action thresholds for codling moth (Lepidoptera: Tortricidae) with pear ester- and codlemone-baited traps in apple orchards treated with sex pheromone mating disruption, 739
Kuenen LPS, Brandl D, Rice RE. Modification of assembly of Pherocon® IC traps speeds trap liner changes and reduces in-field preparation time, 117
Kuhlmann U. See Hemachandra *et al.*, 182
Laffin RD, Dosdall LM, Sperling FAH. Population structure and phylogenetic relationships of *Ceutorhynchus neglectus* (Coleoptera: Curculionidae), 672
Larrivée M. See Pearce *et al.*, 233
Le Lannic J. See Biron *et al.*, 163
Lessard J-P, Buddle CM. The effects of urbanization on ant assemblages (Hymenoptera: Formicidae) associated with the Molson Nature Reserve, Quebec, 215
Lewis TM. See Horton *et al.*, 328
Li C. See Barclay *et al.*, 566
Light DM. See Knight and Light, 728
See Knight and Light, 739
Lindquist EE. See Magowski *et al.*, 648
Macedo PA. See Macedo *et al.*, 497
Macedo TB. See Nansen *et al.*, 428
Macedo TB, Macedo PA, Peterson RKD, Weaver DK, Morrill WL. Rearing the wheat stem sawfly on an artificial diet, 497
Mackauer M. See Danyk *et al.*, 356
Magnusson AK. See Febria *et al.*, 723
Magowski WL, Lindquist EE, Moser JC. *Giselia arizonica*, a new genus and species of mite (Acari: Tarsonemidae) associated with bark beetles of the genus *Pseudopityophthorus* (Coleoptera: Scolytidae) in North America, 648
Maier CT. First North American records of *Batrachedra pinicolella* (Lepidoptera: Batrachedridae), a Palearctic needleminer of spruces, 188
Majka CG. New records of Melyridae (Coleoptera) from the Maritime Provinces of Canada, 325
The Palearctic species *Bembidion femoratum* and *Amara communis* (Coleoptera: Carabidae): new records and notes on modes of introduction to North America, 532
The linden bark borer (Lepidoptera: Agonoxenidae) infesting European linden in Nova Scotia, 620
Malvar RA. See Butrón *et al.*, 350
Malý V. See Howden and Malý, 274
Marshall SA, Paiero SM, Buck M. Buprestid sampling at nests of *Cerceris fumipennis* (Hymenoptera, Crabronidae) in southern Ontario: the first Canadian records of three buprestids (Coleoptera, Buprestidae), 416
Mason PG. See Hemachandra *et al.*, 182
Matheson FO. See Wylie *et al.*, 91
McClay AS. See Mitchell *et al.*, 129
McKee J. See Pearce *et al.*, 233
McKelvey SR. See Fettig *et al.*, 748
McKenney D. See Pearce *et al.*, 233
Meléndez-Ramírez V. See Pinkus-Rendon *et al.*, 441
Meloche F, Rhainds M, Roy M, Brodeur J. Distribution of western and northern corn rootworms (Coleoptera: Chrysomelidae) in Quebec, Canada, 226
Mercado A. See Buddle *et al.*, 120
Miller JR. See Stelinski *et al.*, 98

© 2005 Entomological Society of Canada

Mitchell A, McClay AS, Pohl GR, Sperling FAH. PCR-based methods for identification of two *Eteobalea* species (Lepidoptera: Cosmopterigidae) used as biocontrol agents of weedy *Linaria* species (Scrophulariaceae), 129
Mizuki I, Osawa N, Tsutsumi T. Thrips (Thysanoptera: Thripidae) on the flowers of a dioecious plant, *Dioscorea japonica* (Dioscoreaceae), 712
Moraza ML. A new genus and species of Epicriidae (Acari, Mesostigmata) from eastern North America, 539
Morewood WD, Hoover K, Neiner PR, Sellmer JC. Complete development of *Anoplophora glabripennis* (Coleoptera: Cerambycidae) in northern red oak trees, 376
Morrill WL. See Nansen *et al.*, 202
See Macedo *et al.*, 497
See Nansen *et al.*, 602
Moser JC. See Magowski *et al.*, 648
Mostafa AM. See Wise *et al.*, 598
Muller F. See Gibson *et al.*, 381
Nansen C, Weaver DK, Sing SE, Runyon JB, Morrill WL, Grieshop MJ, Shannon CL, Johnson ML. Within-field spatial distribution of *Cephus cinctus* (Hymenoptera: Cephidae) larvae in Montana wheat fields, 202
Nansen C, Macedo TB, Weaver DK, Peterson RKD. Spatiotemporal distributions of wheat stem sawfly eggs and larvae in dryland wheat fields, 428
Nansen C, Payton ME, Runyon JB, Weaver DK, Morrill WL, Sing SE. Preharvest sampling plan for larvae of the wheat stem sawfly, *Cephus cinctus* (Hymenoptera: Cephidae), in winter wheat fields, 602
Nealis VG. Diapause and voltinism in western and 2-year-cycle spruce budworms (Lepidoptera: Tortricidae) and their hybrid progeny, 584
Neiner PR. See Morewood *et al.*, 376
Nénon JP. See Biron *et al.*, 163
Neven LG. See Horton *et al.*, 328
Nieto Nafría JM. See García Prieto and Nieto Nafría, 49
Nieves-Aldrey JL. Notes on the Neotropical species of *Synergus* (Hymenoptera, Cynipidae), with description of a new species from Colombia, 501
Nobuyoshi O. See Xinhua *et al.*, 83
Novelo-Gutiérrez R, Gómez-Anaya JA. Description of the larva of *Telebasis digiticollis* (Odonata: Zygoptera: Coenagrionidae), 61
Olfert O. See Cárcamo *et al.*, 476
Ordás A. See Butrón *et al.*, 350
Ordás B. See Butrón *et al.*, 350
Osawa N. See Mizuki *et al.*, 712
Owen DR, Wood DL, Parmeter JR. Association between *Dendroctonus valens* and black stain root disease on ponderosa pine in the Sierra Nevada of California, 367
Paiero SM. See Marshall *et al.*, 416
Parmeter JR. See Owen *et al.*, 367
Parra-Tabla V. See Pinkus-Rendon *et al.*, 441
Payton ME. See Nansen *et al.*, 602
Pearce JL, Schuurman D, Barber KN, Larrivée M, Venier LA, McKee J, McKenney D. Pitfall trap designs to maximize invertebrate captures and minimize captures of nontarget vertebrates, 233
Peck SB, Cook J. The genus *Eucatops* of Costa Rica (Coleoptera: Leiodidae, Cholevinae, Eucatopini), 283
Pelletier G. See Klimaszewski *et al.*, 1
Peterson RKD. See Nansen *et al.*, 428
See Macedo *et al.*, 497
Pinkus-Rendon MA, Parra-Tabla V, Meléndez-Ramírez V. Floral resource use and interactions between *Apis mellifera* and native bees in cucurbit crops in Yucatán, México, 441
Pinto JD. See Di Giulio *et al.*, 657
Pohl GR. See Mitchell *et al.*, 129
Price J. See Klimaszewski *et al.*, 1
Qi B. A taxonomic study of Isometopinae from China, including *Myiomma qinlingensis* sp. nov. (Hemiptera: Heteroptera: Miridae), 509
Ranius T. See Wikars *et al.*, 304
Revilla P. See Butrón *et al.*, 350
Rhainds M. See Meloche *et al.*, 226
Rice RE. See Kuenen *et al.*, 117

© 2005 Entomological Society of Canada

Roe AD. See Du *et al.*, 685
Roques A. See Rouault *et al.*, 450
Rouault G, Cantini R, Battisti A, Roques A. Geographic distribution and ecology of two species of *Orsillus* (Hemiptera: Lygaeidae) associated with cones of native and introduced Cupressaceae in Europe and the Mediterranean Basin, 450
Roy M. See Meloche *et al.*, 226
Runyon JB. See Nansen *et al.*, 202
See Nansen *et al.*, 602
Sabahi Q, Talebi K. Demographic studies of *Oomyzus incertus* (Hymenoptera: Eulophidae), a larval parasitoid of *Hypera postica* (Coleoptera: Curculionidae), treated with four organophosphorous insecticides, 488
Sackett TE. See Buddle *et al.*, 120
Sahlin E. See Wikars *et al.*, 304
Sandoya G. See Butrón *et al.*, 350
Sarlis GP. See Kavallieratos *et al.*, 516
Schaefer PW. See Gries *et al.*, 471
Schuurman D. See Pearce *et al.*, 233
Selby RD. See Buddle *et al.*, 120
Sellmer JC. See Morewood *et al.*, 376
Shannon CL. See Nansen *et al.*, 202
Shanower TG. Occurrence of *Sitodiplosis mosellana* (Diptera: Cecidomyiidae) and its parasitoid, *Macroglenes penetrans* (Hymenoptera: Platygasteridae), in northeastern Montana, 753
Shore T. See Barclay *et al.*, 566
Short BD, Bergh JC. Separation of three common hover fly predators of woolly apple aphid based on the exochorionic sculpturing of eggs, 67
Sing SE. See Nansen *et al.*, 202
See Nansen *et al.*, 602
Smith SL. See Fettig *et al.*, 748
Sperling FAH. See Mitchell *et al.*, 129
See Laffin *et al.*, 672
See Du *et al.*, 685
Starý P. See Kavallieratos *et al.*, 516
Stelinski LL, Miller JR, Gut LJ. Captures of two leafroller moth species (Lepidoptera: Tortricidae) in traps baited with varying dosages of pheromone lures or commercial mating-disruption dispensers in untreated and pheromone-treated orchard plots, 98
Storer AJ. See Erbilgin *et al.*, 337
Sugaya H. Revision of the East Asian *Plagiophorus fujiyamai* species group (Coleoptera: Staphylinidae, Pselaphinae, Cyathigerini), 138
Revision of the East Asian *Plagiophorus amygdalinus* species group (Coleoptera: Staphylinidae, Pselaphinae, Cyathigerini), 149
Sweeney J. See Klimaszewski *et al.*, 1
Talebi K. See Sabahi and Talebi, 488
Taylor S. See Barclay *et al.*, 566
Teichert S, Bondrup-Nielsen S. Effect of spatial scale on habitat use of *Bolitotherus cornutus* (Coleoptera: Tenebrionidae), 192
Thomas JC. See Williams and Thomas, 158
Tomanović Ž. See Kavallieratos *et al.*, 516
Trimble RM, El-Sayed AM. Potential of ethyl (2*E*,4*Z*)-2,4-decadienoate for monitoring activity of codling moth (Lepidoptera: Tortricidae) in eastern North American apple orchards, 110
Tsutsumi T. See Mizuki *et al.*, 712
Uddin MJ. See Wylie *et al.*, 91
Ulmer B. See Gibson *et al.*, 381
Ulyshen MD, Hanula JL, Horn S. Using Malaise traps to sample ground beetles (Coleoptera: Carabidae), 251
Varady-Szabo H. See Buddle *et al.*, 120
Vencl FV. See Xinhua *et al.*, 83
Venier LA. See Pearce *et al.*, 233
von Ellenrieder N. See Garrison and von Ellenrieder, 169
Weaver DK. See Nansen *et al.*, 202
See Nansen *et al.*, 428
See Macedo *et al.*, 497
See Nansen *et al.*, 602
White NDG. See Jian *et al.*, 71

© 2005 Entomological Society of Canada

Wikars L-O, Sahlin E, Ranius T. A comparison of three methods to estimate species richness of saproxylic beetles (Coleoptera) in logs and high stumps of Norway spruce, 304
Williams DD. See Febria *et al.*, 723
Williams PH, Thomas JC. A bumblebee new to the New World: *Bombus distinguendus* (Hymenoptera: Apidae), 158
Wise IL, Elliott BG, Mostafa AM. Seasonal occurrence and within-field distribution of *Lygus lineolaris* on buckwheat in Manitoba, 598
Wood DL. See Erbilgin *et al.*, 337
See Owen *et al.*, 367
Wylie HG, Matheson FO, Uddin MJ, Holliday NJ. Release and establishment studies in Manitoba, Canada, of *Aphidius smithi* (Hymenoptera: Aphidiidae), a parasitoid of *Acyrthosiphon pisum* (Hemiptera: Aphididae), 91
Xinhua F, Nobuyoshi O, Vencl FV, Chaoliang L. Structure, behavior, and the life cycle of an aquatic firefly, *Luciola substriata*, in China, 83
Yoo HJS. See Gries *et al.*, 471
Zeran RM. See Buddle et al., 120
Zhang Y. See Dai *et al.*, 404
Zhang Y, Huang M. Taxonomic study of the Oriental genus *Amurta* (Hemiptera: Cicadellidae: Typhlocybinae: Typhlocybini), including *A. bimaculata* sp. nov. and *A. upunctata* sp. nov., 410
Žikić V. See Kavallieratos *et al.*, 516

© 2005 Entomological Society of Canada